

Endo-fullerenes and Doped Diamond Nanocrystallite based Solid-State Qubits

Seongjun Park¹, D. Srivastava², K. Cho¹

¹Multiscale Simulation Lab., Stanford University

²Computational Nanotechnology, NASA Ames Research Center

* Supported by NASA Ames DDF Award to Srivastava and Cho

Background: Kane's Model of Solid-State Quantum Computers

- Solid state quantum computer: Arrays of phosphorus atom in bulk Si, controlled by electronic gates using hyperfine interactions [1]
- Problem: Uniform arrays of individual dopant atoms in bulk Si are experimentally difficult !




1. Kane, B.E., *Nature*, 393, p.133 (1998)

Mar. 13 2001

MSL Stanford & NASA Ames


2/11

Solution: Use Encapsulated Atoms as Qubits !



Proposal: Arrays of "encapsulated" atoms (with 1/2 nuclear spin – qubits) will be easy to fabricate as compared to the arrays of the similar bare atoms.

Example: ¹H encapsulated in C₃₆



Electronic charge density shows a weak meta-stable state of ¹H at the center of C₃₆

Suitable Solid-state Qubits Identified:

- ¹H encapsulated in a C₃₆D₂₀ fullerene
- ³¹P encapsulated in a diamond nanocrystallite

Mar. 13 2001


MSL Stanford & NASA Ames

3/11

Example: ¹H Encapsulated in C₃₆

- Center is meta-stable site.
- ¹H strongly prefers to make a bond with a carbon atom

	Formation Energy (eV)
Carbon A	-1.28
Carbon B	-1.54
Carbon C	-1.40
Center	-0.46




Mar. 13 2001

MSL Stanford & NASA Ames

4/11

Reactivity Control for ¹H: C₂₀D₂₀

- ¹H prefers to make a bond with C atom within fullerene.
→ Reduce the chemical reactivity
- sp³ hybrid will reduce the electron density of inner surface.
→ Hydrogenation
- Hexagon has lower diffusion barrier than pentagon.
→ Non-hexagon structure is preferred.
- As a conclusion, we examined C₂₀D₂₀




Mar. 13 2001

MSL Stanford & NASA Ames

5/11

¹H Encapsulated in C₂₀D₂₀

- Energy changes as moving ¹H from the center to the wall of C₂₀D₂₀



- Symmetry sites: Carbon, Bond, and Pentagon
- Center is the most stable site.
- No meta-stable site
- Diffusion Barrier ~ 1.17 eV


Mar. 13 2001

MSL Stanford & NASA Ames

6/11

Charge Density of ^1H Encapsulated in $\text{C}_{20}\text{D}_{20}$

- Some of electron wave of ^1H is found out of $\text{C}_{20}\text{D}_{20}$



Mar. 13 2001

MSL, Stanford & NASA Ames

7/1


Model 2: Doped Diamond Nanocrystallite Step1: ^{31}P Encapsulation in C_{60}

- Center is the most stable site for ^{31}P
- Diffusion barrier to move to a wall C atom is about 0.35 eV

^{31}P in C_{60}

	Formation Energy (eV)
(6,6) Bond	-0.81
Center	-0.99

- Implantation of ^{31}P in a C_{60} fullerenes has been experimentally done!



Knappe, C. et al. *Molecular Physics* 95, 999-1004 (1998).


Mar. 13 2001

MSL, Stanford & NASA Ames

8/1

Steps 2-3: Convert to a ^{31}P Encapsulation in a Diamond Nanocrystallite

- Step2: grow bucky-onion layers with ^{31}P @ C_{60} fullerenes as seed material
- Step3: e-beam irradiation of the bucky-onion layers converts the core shells into ^{31}P doped in a diamond nanocrystallite



Benhart, F. and Ajayan, P.M., *Nature*, 382, p. 443 (1996)


Mar. 13 2001

MSL, Stanford & NASA Ames

9/1

^{31}P doping in Diamond or Silicon

- Energy difference between ^{31}P at substitutional site, and two types of interstitial site of compressed diamond (0, 20, and 50 GPa) and bulk Si



- Substitutional site is more stable than interstitial site.


- Diamond has bigger energy differences than Si.

Mar. 13 2001

MSL, Stanford & NASA Ames

10/1

Charge Density of ^{31}P in Diamond or Silicon



^{31}P in Diamond

^{31}P in Si

Mar. 13 2001

MSL, Stanford & NASA Ames

11/1

^{31}P in a Diamond Nanocrystallite as a Qubit (Comments)

- ^{31}P at a substitutional site in a diamond nanocrystallite is more stable (by ~15 eV) than a ^{31}P at an interstitial site - no diffusion
- Spin-orbit interaction on a qubit is 8 times stronger than Kane's model
- In principle any host lattice can be chosen to tune the neighboring qubit interactions
- Experimentally feasible fabrication pathway for these solid-state qubits (Steps 1-3) exist

Mar. 13 2001

MSL, Stanford & NASA Ames

12/1